

STK16 Human

Description: STK16 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 329 amino acids (1-305 a.a) and having a molecular mass of 37.2kDa. STK16 is fused to a 24 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #: PKPS-039

For research use only.

Synonyms: Serine/threonine-protein kinase 16, Myristoylated and palmitoylated serine/threonine-protein kinase, MPSK, Protein kinase PKL12, TGF-beta-stimulated factor 1, TSF-1, Tyrosine-protein kinase STK16, hPSK, STK16, MPSK1, PKL12, TSF1, KRCT.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MGSHMGHALC VCSRGTVID
NKRYLFIQKL GEGGFSYVDL VEGLDHGHFY ALKRILCHEQ QDREEAQREA DMHRLFNHPN
ILRLVAYCLR ERGAKHEAWL LLPFFKRGTLL WNEIERLKDK GNFLTEDQIL WLLLGICRGL
EAIHAKGYAH RDLKPTNILL GDEGQPVLM D LGSMNQACIH VEGSRQALTL QDWAAQRCTI
SYRAPELFSV QS

Purity: Greater than 90.0% as determined by SDS-PAGE.

Formulation:

STK16 protein solution (1mg/ml) containing 20mM Tris-HCl buffer (pH8.0), 20% glycerol, 0.1M NaCl and 1mM DTT.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Serine/threonine-protein kinase 16 (STK16) is a membrane-associated protein kinase which phosphorylates on serine and threonine residues. STK16 is involved in secretory vesicle trafficking or intracellular signaling. Furthermore, the STK16 protein may have a role in regulating stromal-epithelial interactions which occur during ductal morphogenesis in the mammary gland. STK16 can autophosphorylate on Tyr residue; it is however unclear whether STK16 has tyrosine-protein kinase toward other proteins. STK16 may also be involved in TGF-beta signaling.

To place an order, please Click [HERE](#).