

FN3K Human

Description:FN3K Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 332 amino acids (1-309 a.a) and having a molecular mass of 37kDa.FN3K is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #:PKPS-056

For research use only.

Synonyms:Fructosamine-3-kinase, FN3K.

Source:Escherichia Coli.

Physical Appearance:Sterile filtered colorless solution.

Amino Acid Sequence:MGSSHHHHH SSGLVPRGSH MGSMEQLLRA ELRTATLRAF
GGPGAGCISE GRAYDTDAGP VFKVNRRTQ ARQMFEGEVA SLEALRSTGL VRVPRPMKVI
DLPGGGAFFV MEHLKMKSL SQA SKLGEQM ADLHLYNQKL REKLKEEENT VGRRGEGAEP
QYVDKFGFHT VTCCGFIPQV NEWQDDWPTF FARHRLQAQL DLIEKDYADR EARELWSRLQ
VKIPDLFCGL EI

Purity:Greater than 85.0% as determined by SDS-PAGE.

Formulation:

FN3K protein solution (0.25mg/ml) containing 20mM Tris-HCl buffer (pH 8.0), 0.15M NaCl, 20% glycerol and 1mM DTT.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. They may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Fructosamine 3 Kinase (FN3K) catalyzes the phosphorylation of fructosamines which may result in deglycation, the non-enzymatic reaction of glucose with primary amines followed by Amadori re-arrangement. Phosphorylation of fructosamines instigates metabolism of the modified amine and brings about the de-glycation of fructoselysine and of glycated proteins. A high concentration of glucose may affect non-enzymatic oxidation of proteins by reaction of glucose and lysine residues (glycation). Fructosamines, the proteins altered in this way, are less active or functional.

To place an order, please [Click HERE](#).