www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

## MMACHC Human

Description: MMACHC Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 306 amino acids (1-282 a.a) and having a molecular mass of 34kDa.MMACHC is fused to a 24 amino acid His-tag at N-terminus & Durified by proprietary chromatographic techniques.

Catalog #:PRPS-1126

For research use only.

Synonyms: Methylmalonic aciduria and homocystinuria type C protein, MMACHC, cblC, RP11-291L19.3.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MGSHMEPKVA ELKQKIEDTL CPFGFEVYPF QVAWYNELLP PAFHLPLPGP TLAFLVLSTP AMFDRALKPF LQSCHLRMLT DPVDQCVAYH LGRVRESLPE LQIEIIADYE VHPNRRPKIL AQTAAHVAGA AYYYQRQDVE ADPWGNQRIS GVCIHPRFGG WFAIRGVVLL PGIEVPDLPP RKPHDCVPTR ADRIALLEGF NFHWRDWTYR DA

Purity: Greater than 90.0% as determined by SDS-PAGE.

#### Formulation:

MMACHC protein solution (0.5mg/ml) containing 20mM Tris-HCl buffer (pH 8.0), 0.15M NaCl, 10% glycerol and 1mM DTT.

### Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

# Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

## Introduction:

Though the exact role of MMACHC is unknown, its C-terminal region shows similarity to TonB, which is a bacterial protein involved in energy transduction for cobalamin (vitamin B12) uptake. Therefore, it is suggested that MMACHC may have a role in the binding and intracellular trafficking of cobalamin. MMACHC mutations are linked with methylmalonic aciduria and homocystinuria type cblC. MMACHC is widely expressed, it is expressed at higher level in the fetal liver, however it is also expressed in the spleen, lymph node, thymus and bone marrow. MMACHC is weakly or not expressed in the peripheral blood leukocytes.

To place an order, please Click HERE.





