

## WBP2 Human

**Description:**WBP2 Human Recombinant produced in E. coli is a single polypeptide chain containing 284 amino acids (1-261) and having a molecular mass of 30.5 kDa.WBP2 is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #:PRPS-1215

**Synonyms:**WW domain binding protein 2, WBP-2.

For research use only.

**Source:**E.coli.

**Physical Appearance:**Sterile Filtered colorless solution.

**Amino Acid Sequence:**MGSSHHHHHH SSGLVPRGSH MGSMLNKNH SEGGGVIVNN  
TESILMSYDH VELTFNDMKN VPEAFKGTCK GTVYLTPYRV IFLSKGKDAM QSFMMPFYLM  
KDCEIKQPVF GANYIKGTVK AEAGGGWEGS ASYKLTFTAG GAIEFGQRML QVASQASRGE  
VPSGAYGYSY MPSPAYVYPP PVANGMYPYCP PGYPYPPPPP EFYPGPPMMD GAMGYVQPPP  
PPYPGPMEPP VS

**Purity:**Greater than 90% as determined by SDS-PAGE.

### Formulation:

The WBP2 solution (0.5mg/1ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.1M NaCl, 1mM DTT and 20% glycerol.

### Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.

### Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

### Introduction:

WW domain-binding protein 2 (WBP2) is a steroid hormone receptor coactivator, which contains 1 GRAM domain. The globular WW domain is comprised of 38-40 semiconserved amino acids shared by proteins of various functions including structural, regulatory, and signaling proteins. This domain is involved in mediating protein-protein interactions through the binding of polyproline ligands. WBP2 binds to the WW domain of Yes kinase-associated protein(YAP1) via its PY motifs.

**To place an order, please [Click HERE](#).**