

BCL2L10 Human

Description:BCL2L10 Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 195 amino acids (1-172 a.a.) and having a molecular mass of 21.8kDa.BCL2L10 is fused to a 23 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #:PRPS-1328

For research use only.

Synonyms:Bcl-2-like protein 10, Bcl2-L-10, Anti-apoptotic protein NrH, Apoptosis regulator Bcl-B, BCL2L10, BCLB, Boo, Diva, BCL-B.

Source:E.coli.

Physical Appearance:Sterile Filtered colorless solution.

Amino Acid Sequence:MGSSHHHHHH SSGLVPRGSH MGSMVDQLRE RTTMADPLRE
RTELLADYL GYCAREPGTP EPAPSTPEAA VLRSAAARLR QIHRFFSAY LGYPGNRFEL
VALMADSVLS DSPGPTWGRV VTLVTFAGTL LERGPLVTAR WKKWGFQPRL KEQEGDVARD
CQRLVALLSS RLMGQHRAWL QAQGGWDGFC HFFRT.

Purity:Greater than 90% as determined by SDS-PAGE.

Formulation:

BCL2L10 protein solution (0.5mg/ml) containing 20mM Tris-HCl buffer (pH 8.0), 0.15M NaCl, 20% glycerol and 1mM DTT.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA).Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

BCL2 Like 10 (BCL2L10) is a member of the BCL-2 protein family. BCL2 family members form hetero or homodimers and serve as anti or pro-apoptotic regulators which are involved in a wide variety of cellular activities. BCL2L10 includes the conserved BH4, BH1 and BH2 domains. BCL2L10 interacts with other members of BCL2 protein family including BCL2, BCL2L1/BCL-X(L), and BAX. BCL2L10 overexpression suppresses cell apoptosis possibly through the prevention of cytochrome C release from the mitochondria, and consequently activating caspase-3 activation.

To place an order, please [Click HERE](#).