

DEDD Human

Description:DEDD Human Recombinant produced in E.Coli is a single, non-glycosylated polypeptide chain containing 338 amino acids (1-318a.a) and having a molecular mass of 38.9kDa. DEDD is fused to a 20 amino acid His-tag at N-terminus & purified by proprietary chromatographic techniques.

Catalog #:PRPS-1356

For research use only.

Synonyms:CASP8IP1, DEDD1, DEFT, FLDED1, KE05, DEDPro1, Death effector domain-containing testicular molecule.

Source:Escherichia Coli.

Physical Appearance:Sterile Filtered colorless solution.

Amino Acid Sequence:MGSSHHHHHH SSGLVPRGSH MAGLKRRASQ VWPEEHGEQE
HGLYSLHRMF DIVGTHLTHR DVRVLSFLFV DVIDDHERGL IRNGRDFLA LERQGRCDDES
NFRQVLQLLR IITRHDLLPY VTLKRRRAVC PDLVDKYLEE TSIRYVTPRA LSDPEPRPPQ
PSKTVPPHYP VVCCPTSGPQ MCSKRPARGR ATLGSRKRR KSVTPDPKEK QTCDIRLRVR
AEYCQHETAL QG

Purity:Greater than 85.0% as determined by SDS-PAGE.

Formulation:

DEDD protein solution (0.5mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 0.4M Urea and 10% glycerol.

Stability:

A2LD1 should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

Death effector domain-containing protein (DEDD) is a cytoplasmic protein. The cell death activity of DEDD relates to its nuclear localization. DEDD translocates to the nucleus during CD95-mediated apoptosis, there it localizes to nucleoli-like structures, activates caspase-6 and particularly inhibits RNA polymerase I-dependent transcription. DEDD is usually expressed in a variety of tissues, and found in the highest levels in the testis. Overexpression of DEDD was shown to induce weak apoptosis.

To place an order, please [Click HERE](#).