www.neobiolab.com info@neobiolab.com 888.754.5670, +1 617.500.7103 United States 0800.088.5164, +44 020.8123.1558 United Kingdom

S100A14 Human

Description: \$100A14 Human Recombinant fused with a 20 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 124 amino acids (1-104 a.a.) and having a molecular mass of 13.8kDa. The S100A14 is purified by proprietary chromatographic techniques.

Catalog #:PRPS-161

For research use only.

Synonyms: Protein S100-A14, S100 calcium-binding protein A14, S114, S100A14, S100A15, BCMP84.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MGQCRSANAE DAQEFSDVER AIETLIKNFH QYSVEGGKET LTPSELRDLV TQQLPHLMPS NCGLEEKIAN LGSCNDSKLE FRSFWELIGE AAKSVKLERP VRGH.

Purity: Greater than 95.0% as determined by SDS-PAGE.

Formulation:

The S100A14 solution (1 mg/ml) contains 20mM Tris-HCl buffer (pH 8.0), 10% glycerol and 2mM DTT.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks. Store, frozen at -20°C for longer periods of time. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Avoid multiple freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

S100A14 belongs to a subfamily of proteins related by EF-hand Ca2+ binding protein superfamily. The extracellular functions of the S100 family include the ability to boost neurite outgrowth, involvement in inflammation, and motility of tumor cells. S100A14 contains 2 EF-hand Ca2+-binding domains, a myristoylation motif, a glycosylation site, and a number of potential protein kinase phosphorylation sites. S100A14 is expressed at high levels in the colon and at moderate levels in the thymus, kidney, liver, small intestine, and lung.

To place an order, please Click HERE.





