

RORC Human

Description:RAR-Related Orphan Receptor C Human Recombinant produced in E.Coli is a full length protein consisting of 497 amino acids having a molecular weight of 55.8kDa and fused with 5.5kDa amino-terminal His-Flag tag.RORC is purified by proprietary chromatographic techniques.

Catalog #:PRPS-213

Synonyms:Nuclear receptor ROR-gamma, Nuclear receptor RZR-gamma, Nuclear receptor subfamily 1 group F member 3, Retinoid-related orphan receptor-gamma, RORC, NR1F3, RORG, RZRG, TOR, RZR-GAMMA.

For research use only.

Source:Escherichia Coli.

Physical Appearance:Sterile filtered colorless solution.

Amino Acid Sequence:

MSYHHHHHHHDYDIPTTDYKDDDDKDYKDDDDKENLYFQGEFMRTQIEVIPCKICGDKSSGIHYG
VITCEGCKGFFRRSQRCNAAYSCTRQQNCPIDRTSRNRCQHCRLLQKCLALGMSRDAVKFGRMS
KKQRDSLHAEVQKQLQQRQQQQEPVVKTPPAGAQQADTLTYTLGLPDGQLPLGSSPDLPEAS
ACPPGLLKASGSGPSYSNNLAKAGLNGASCHLEYSPEERGKAEGRESFYSTGSQQLTPDRCLRFE

Purity:Greater than 80.0% as determined by SDS-PAGE.

Formulation:

RORC protein is supplied in 50mM Tris, 150mM NaCl and 10% Glycerol, pH 7.5.

Stability:

Store at 4°C if entire vial will be used within 2-4 weeks.Store, frozen at -20°C for longer periods of time.Please avoid freeze thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. They may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

RORC is a DNA-binding transcription factor which belongs to the NR1 subfamily of nuclear hormone receptors. The specific functions of the RORC protein are not known; nevertheless, studies of a similar gene in mice have shown that the RORC gene may be vital for lymphoid organogenesis and may have an imperative regulatory role in thymopoiesis. Furthermore, studies in mice suggest that RORC may inhibit the expression of Fas ligand and IL2. RORC may be a possible nuclear receptor for hydroxycholesterols, the binding of which strongly promotes coactivators recruitment. RORC is Crucial for thymopoiesis and the development of several secondary lymphoid tissues, including lymph nodes. RORC is also involved in lineage specification of uncommitted CD4(+) T helper cells into Th17 cells. In addition, RORCs regulate the expression of several components of the circadian clock.

To place an order, please [Click HERE](#).