

CDCA8 Human

Description: CDCA8 Human Recombinant fused with a 21 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 283 amino acids (19-280a.a.) and having a molecular mass of 31.6kDa. The CDCA8 is purified by proprietary chromatographic techniques.

Catalog #: PRPS-229

For research use only.

Synonyms: Cell division cycle associated 8, BOR, hDasraB, FLJ12042, MESRGP, Pluripotent embryonic stem cell-related gene 3 protein, PESCRG3, BOREALIN.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MRKLASFLKD FDREVEIRIK
QIESDRQNLL KEVDNLYNIE ILRLPKALRE MNWLDYFALG GNKQALEEAA TADLDITEIN
KLTAELAIQTP LKSAKTRKVI QVDEMIVEEE EEEENERKNL QTARVKRCPP SKKRTQSIQG
KGKGKRSSRA NTVTPAVGRLEVSMVKPTPG LTPRFDSRVF KTPGLRTPAA GERIYNISGN
GSPLADSKEI FLT

Purity: Greater than 90.0% as determined by SDS-PAGE.

Formulation:

The CDCA8 solution (0.25mg/1ml) contains 20mM Tris-HCl buffer (pH 8.0), 50% glycerol, 1mM DTT, 1mM EDTA and 0.2M NaCl.

Stability:

CDCA8 should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

CDCA8 is a component of the chromosomal passenger complex (CPC). The CPC complex has crucial functions at the centromere in ensuring accurate chromosome placement and isolation and is obligatory for chromatin-induced microtubule stabilization and spindle assembly. In the complex, CDCA8 protein is needed to direct the CPC to centromeric DNA. Removal of CDCA8 delays mitotic progression and causes kinetochore-spindle to mis-attach.

To place an order, please [Click HERE](#).