

TCP1 Human

Description: TCP1 Human Recombinant fused with a 20 amino acid His tag at N-terminus produced in E.Coli is a single, non-glycosylated, polypeptide chain containing 576 amino acids (1-556 a.a.) and having a molecular mass of 62.5kDa. The TCP1 is purified by proprietary chromatographic techniques.

Catalog #:PRPS-283

For research use only.

Synonyms: T-complex protein 1 subunit alpha, TCP-1-alpha, CCT-alpha, TCP1, CCT1, CCTA, D6S230E.

Source: Escherichia Coli.

Physical Appearance: Sterile Filtered colorless solution.

Amino Acid Sequence: MGSSHHHHHH SSGLVPRGSH MEGPLSVFGD RSTGETIRSQ
NVMAAASIAN IVKSSLGPVG LDKMLVDDIG DVTITNDGAT ILKLLEVEHP AAKVLCCELAD
LQDKEVGDGT TSVVIAAEL LKNADELVKQ KIHPTSVISG YRLACKEAVR YINENLIVNT
DELGRDCLIN AAKTSMSSKI IGINGDFFAN MVVDAVLAIK YTDIRGQPRY PVNSVNILKA
HGRSQMESML IS

Purity: Greater than 80.0% as determined by SDS-PAGE.

Formulation:

The TCP1 solution (0.5 mg/ml) contains 20mM Tris-HCl Buffer (pH 8.0), 1mM DTT, 0.1mM PMSF and 10% Glycerol.

Stability:

TCP1 should be stored desiccated below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

Usage:

NeoBiolab's products are furnished for LABORATORY RESEARCH USE ONLY. The product may not be used as drugs, agricultural or pesticidal products, food additives or household chemicals.

Introduction:

TCP1 is a molecular chaperone that is a member of the chaperonin containing TCP1 complex (CCT), also known as the TCP1 ring complex (TRiC). This complex consists of 2 identical stacked rings, each containing eight different proteins. Unfolded polypeptides penetrate the central cavity of the complex and are folded in an ATP-dependent manner. The TCP1 protein is found in the cytosol as a subunit of a hetero-oligomeric chaperone. TCP1 has a significant function in maintaining cellular homeostasis by assisting the folding of many proteins such as the cytoskeletal components actin and tubulin.

To place an order, please [Click HERE](#).